

The effect of turbulent intermittency on the deflagration to detonation transition in SN Ia explosions

Liubin Pan, J. Craig Wheeler and John Scalapino

Astronomy Department, University of Texas at Austin

panlb@astro.as.utexas.edu

ABSTRACT

We examine the effects of turbulent intermittency on the deflagration to detonation transition (DDT) in Type Ia supernovae. The Zel'dovich mechanism for DDT requires the formation of a nearly isothermal region of mixed ash and fuel that is larger than a critical size. We primarily consider the hypothesis by Khokhlov et al. and Niemeyer and Woosley that the nearly isothermal, mixed region is produced when the flame makes the transition to the distributed regime. We use two models for the distribution of the turbulent velocity fluctuations to estimate the probability as a function of the density in the exploding white dwarf that a given region of critical size is in the distributed regime due to strong local turbulent stretching of the flame structure. We also estimate lower limits on the number of such regions as a function of density. We find that the distributed regime, and hence perhaps DDT, occurs in a local region of critical size at a density at least a factor of 2 – 3 larger than predicted for mean conditions that neglect intermittency. This factor brings the transition density to be much larger than the empirical value from observations in most situations. We also consider the intermittency effect on the more stringent conditions for DDT by Lisewski et al. and Woosley. We find that a turbulent velocity of 10^8 cm/s in a region of size 10^6 cm, required by Lisewski et al., is rare. We expect that intermittency gives a weaker effect on the Woosley model with stronger criterion. The predicted transition density from this criterion remains below 10^7 g/cm³ after accounting for intermittency using our intermittency models.

Subject headings: stars: interiors—supernovae: general—turbulence

1. Introduction

A successful model for Type Ia Supernova (SNIa) explosions is required to produce a deflagration to detonation transition (DDT) by observational constraints. A pure deflagration

model gives exploding kinetic energy lower than observed (Khokhlov 1991; Gamezo et al. 2003; Röpke & Hillebrandt 2005) and pure detonation leads to overproduction of iron group elements and too little intermediate elements (Branch et al. 1982, 1983). The density ρ_{tr} at which the transition occurs determines the amount of the nickel produced (Höflich 1995; Höflich and Khokhlov 1996; Dominguez, Höflich & Straniero 2001). Therefore a prediction of ρ_{tr} , consistent with the observed nickel production, is essential to a DDT theory for SNe Ia .

The mechanism by which the DDT occurs still remains a mystery. The most studied candidate is the Zel’dovich mechanism, which requires the existence of an almost isothermal region of mixed ash and fuel that is larger than a critical size l_c to drive a supersonic shock that is sufficiently strong to sweep over the entire star (Khokhlov et al. 1997, hereafter KOW; Niemeyer and Woosley 1997, hereafter NW). One hypothesis is that a nearly isothermal region is produced by turbulent preconditioning. KOW argued that, to produce an almost isothermal mixture of ash and fuel, the laminar flame must be quenched by turbulent stretching, at least locally. This might allow the cold fuel to mix with the ash both thermally by electron conduction and chemically by diffusivity without being burned. They assumed that the criterion to quench a flame is that the turbulent velocity at the laminar flame thickness must be larger than the laminar flame speed. NW gave a similar argument based on the distributed flame burning regime in turbulent combustion. The criterion for a distributed flame is expressed in terms of the Gibson scale at which the turbulence velocity equals the laminar flame speed. If the Gibson scale is smaller than the laminar flame thickness, turbulent stretching can generate structures within the flame and the flame is in the distributed regime. NW speculated that in this regime flames can be temporally quenched in some regions, which can host the detonation after being homogenized in temperature and composition by turbulent mixing. The criterion for the distributed regime is equivalent to that for flame quenching used by KOW. Both criteria give the same condition on the turbulence intensity for given laminar flame properties (see §2). As the density in the star drops due to the overall expansion, it is easier for turbulence to affect the laminar flame because of the decrease in the flame speed and the increase in flame thickness. With presumed turbulence parameters, the criterion for the turbulence intensity, determined by the robustness of laminar flames disturbed by turbulent motions, translates to a transition density ρ_{tr} for the DDT.

Several uncertainties exist in the simple model given by these two early studies. First, it is not clear whether the criterion used by KOW, equivalent to that for a distributed regime (NW), is sufficient for flame breaking. How, or even if, flames are quenched is still an open question. Second, it is uncertain whether (local) flame quenching is indeed necessary to produce a nearly isothermal region. Finally, later studies by Lisewski, Hillebrandt and

Woosley (2000) (hereafter Lisewski et al 2000; see also Lisewski et al. 2000b) and Woosley (2007) find that entering the distributed regime, while probably a necessary condition, is not sufficient for the DDT to occur. Based on a requirement for turbulent transport to be efficient at producing a shallow temperature and composition gradient around the laminar flame, Lisewski et al. (2000) find that the turbulent velocity at the scale 10^6 cm needed for a detonation is very large, $\sim 10^8$ cm/s. Woosley (2007) claims that the DDT occurs only when the turbulent flame thickness exceeds a critical length scale. We show in §2 that the two criteria, although arising from different physical considerations, are basically equivalent. The corresponding condition is more stringent than that assumed by KOW and NW.

In this paper we examine the effect of turbulent intermittency on the onset of distributed burning that may relate to the DDT. Despite the uncertainties listed above, we will mainly consider the model by KOW and NW and use it to illustrate the potential importance of intermittency in SN Ia explosions. Our calculations can be applied to the criteria by Lisewski et al. (2000) and Woosley (2007) in a straightforward way. A quantitative analysis using their criteria requires data for laminar flame properties and critical length scales at densities below 10^7 g cm $^{-3}$ that are not immediately available (see §2). We give a qualitative discussion of the intermittency effect on their DDT models.

Intermittency is an important concept in turbulence theory. It is characterized by intense local events, e.g., strong stretching at small scales, which occur at a frequency much larger than predicted from a Gaussian distribution (see, e.g., Frisch 1995). The physical origin of intermittency in turbulent flows is the spatial inhomogeneity in the energy dissipation rate: most kinetic energy is viscously dissipated in the finest structures, e.g., vortex tubes, which occupy only a small volume fraction. These rare but intense dissipative structures give rise to a spatially inhomogeneous and intermittent distribution for the turbulent intensity and the stretching rate. Intermittency is shown as broad exponential tails in the probability distribution for the stretching rate or the dissipation rate at small scales (see §3). The tails get broader at smaller scales, meaning that the probability of finding an extreme turbulent stretching rate or intensity increases with decreasing scales.

According to 1D simulation results by KOW and NW, the critical size, l_c , of the isothermal region required for a DDT via the Zel’dovich mechanism is much smaller, especially at large densities, than the expected integral length scale for the buoyancy-driven turbulence in SN Ia explosions. This suggests that only a small flame region with a sufficiently strong local turbulence intensity may be needed to trigger a detonation. Turbulent intermittency, which indicates the existence of regions of small sizes where the turbulent stretching is much larger than the average value over the flow, is therefore expected to have important consequences for DDT. The transition could happen earlier at a higher transition density ρ_{tr} than

predicted by models using the average turbulent intensity. At higher densities, much larger turbulent intensity is required for the DDT, but the rapid decrease of l_c with increasing density makes an earlier DDT possible for two reasons. First, the probability is larger to find regions of smaller sizes l_c with extreme turbulent stretching rate or intensity. Second, there are more regions of smaller size available as candidates to host the detonation. Clearly, the intermittency effect accounts for the intuitive dependence of ρ_{tr} on l_c : the smaller the critical size, the easier it may be for the transition to happen. To what degree the intermittency effect increases ρ_{tr} is the main question we investigate in this paper.

In §2, we review the criteria for the DDT in models by KOW, NW, Lisewski et al. (2007) and Woosley (2007) and formulate a new criterion taking into account the effect of intermittency. We describe two intermittency models by Oboukhov (1962) and Kolmogorov (1962) and by She and Leveque (1994) in §3. Using the intermittency models, we evaluate the transition density from the new criteria in §4. Our results are summarized and discussed in §5.

2. Criteria for the DDT

The criterion used in NW for judging whether a flame is in the distributed regime, which was also assumed to be the condition for the DDT, is to compare the Gibson scale l_G with the laminar flame thickness l_f . The Gibson scale is defined such that $\delta u(l_G) = S_l$ where $\delta u(l)$ is the amplitude of the velocity fluctuations at the scale l (or equivalently the velocity difference over a scale l , i.e., $\delta u(l) = u(l+x) - u(x)$) and S_l is the laminar flame speed¹. If $l_G \gtrsim l_f$, the turbulence cannot internally disturb the flame and the turbulence in effect wrinkles the flame. This is called the flamelet regime. Only when $l_G \lesssim l_f$, can turbulence stretch the flame efficiently to generate structures within the flame and the turbulent combustion enters the distributed regime. The condition $l_G \lesssim l_f$ is equivalent to $\delta u(l_f) \gtrsim \delta u(l_G) = S_l$ since $\delta u(l)$ is an increasing function of the scale l ². The latter, which means that the turbulent

¹Note that NW, accounting for the cellular stabilization effect against instabilities, e.g., the Landau-Darrieus instability, defined l_G as the scale where the turbulent velocity exceeds the effective cellular flame speed. This does not introduce a significant difference in the estimate of l_G since the effective cellular speed is close to the laminar speed and has a very weak dependence on scale; see their Fig. 1.

²This is also equivalent to the diffusivity criterion by Niemeyer and Kerstein (1997) for the onset of distributed regime and flame extinction at Prandtl number larger than unity, which is the case for white dwarfs. Their criterion was motivated by the observation that, at Prandtl number different from unity, two previous criteria proposed for the flamelet breakdown and for the flame quenching, using the ratio of the flow viscous length scale to the flame thickness and the ratio of the viscous timescale to the reaction timescale,

velocity fluctuation $\delta u(l_f)$ at the scale of the flame thickness l_f is larger than the laminar flame speed, is the criterion used in KOW for flame quenching and the DDT.

Following KOW, we introduce a factor of $K \sim 1$ in the criterion to account for the uncertainty in the flame breaking mechanism, i.e., $\delta u(l_f) \geq K S_l$. We will consider two values for K , i.e., $K = 1$ and $K = 8$ (KOW; For $K = 8$, to quench a flame, the Gibson scale has to be $K^3 = 512$ times smaller than the flame width). This criterion can also be written in terms of timescales. Noting that the turbulent stretching timescale, τ_t , at the flame thickness is $\tau_t(l_f) = l_f/\delta u(l_f)$ and that the nuclear reaction timescale, τ_n , is related to the flame speed $\tau_n = l_f/S_l$, the criterion is equivalent to $\tau_t(l_f) < \tau_n/K$, i.e., to break the flame the stretching timescale at the flame thickness must be smaller than the nuclear burning timescale (see Niemeyer and Kerstein 1997).

To apply this criterion, the Kolmogorov (1941) scaling $\delta u(l) = \bar{\epsilon}^{1/3} l^{1/3}$ is usually used to calculate $\delta u(l_f)$ from the turbulent velocity fluctuations at large scales where $\bar{\epsilon}$ is the average dissipation rate in the flow. From this scaling, the criterion can be written as (KOW, NW),

$$\bar{\epsilon}^{1/3} l_f^{1/3} > K S_l \quad (1)$$

or

$$\bar{\epsilon} > K^3 S_l^3 / l_f = K^3 \epsilon_f \quad (2)$$

where ϵ_f is defined as S_l^3/l_f . Although we use the convenient criterion (2) in terms of the dissipation rate in our calculations, the turbulent stretching is more fundamental and we will use the concept of the flame stretching in our discussions.

The laminar flame speed and thickness depend on the chemical composition and the density (Timmes and Woosley 1992, KOW). In Table 1, we list the flame speed, the flame thickness as a function of density for a white dwarf with half carbon and half oxygen, mainly taken from Timmes and Woosley (1992). The laminar speed decreases and the thickness increases quickly with decreasing density ρ , therefore ϵ_f decreases rapidly with decreasing ρ as shown in Table 1. The average dissipation rate is estimated to be $\bar{\epsilon} = U^3/L$ where U and L are the characteristic velocity and length scales of the turbulence, normally set by motions on the large, driving scale. At large scales, the turbulence is driven by the Rayleigh-Taylor instability. The length scale L might be expected to be about the size, R_f , of the flame region, $L \simeq R_f \sim 10^8$ cm and the velocity scale to be about the Rayleigh-Taylor velocity at this scale $U \sim 0.5\sqrt{g_{eff}L} \simeq 10^8$ cm/s where the effective gravity is taken to be $\sim g_{eff} = 5 \times 10^8$ cm/s² (KOW, NW). Khokhlov (1995), however, showed that motions at

respectively, are not equivalent.

scales larger than 10^6 cm freeze out due to the overall expansion of the star. In that case, $L \sim 10^6 - 10^7$ cm and $U \sim 10^7$ cm/s. We will take U and L as parameters. Note that the criterion eq (2) depends on U and L through the dissipation rate. Given the dissipation rate $\bar{\epsilon}$, the critical density below which the inequality (eq 2) is satisfied can be obtained using ϵ_f as a function of ρ in Table 1. For example, if $U \sim 100$ km/s and $L \sim 100$ km, $\bar{\epsilon} \sim 10^{14}$ cm²/s³ and we find that, from interpolation in Table 1, $\bar{\epsilon}$ is larger than ϵ_f at a density less than $\sim 4 \times 10^7$ g/cm³. Therefore criterion (2) predicts a transition density $\rho_{tr} \simeq 4 \times 10^7$ g/cm³ for $K = 1$ (see KOW and NW). If $K = 8$, the predicted transition density is smaller, $\rho_{tr} \sim 1.5 \times 10^7$ g/cm³. In the second line of Table 2, we give the predicted ρ_{tr} for different values of the parameters, which decreases with decreasing dissipation rate $\bar{\epsilon}$. The numbers in parenthesis correspond to $K = 8$.

When using the criterion eq (2), we need to keep in mind that the spatial fluctuations of ϵ (see §3) are completely neglected and the criterion only applies to the overall situation in the combustion flow. We will refer to this criterion as the mean criterion. When the mean criterion is met, the only implication is that the combustion is in the distributed regime in general. Considering the intermittency of turbulence, i.e., the spatially inhomogeneous distribution of the stretching strength, there can be places where the stretching rate is much weaker than the average. These places could still be in the flamelet stage while most other places are in the distributed regime. Or conversely, even if the mean criterion (2) is not satisfied, one cannot exclude the possibility of there existing a region that experiences strong stretching and gets into the distributed regime when most of the structure is still in the flamelet regime. This latter fact is important for the deflagration to detonation transition. The fact that the DDT does not require the entire star to be in the distributed regime but instead only needs a region of size much smaller than the white dwarf radius (see below, KOW), coupled with the intrinsic intermittency, suggests that DDT could occur earlier than predicted by eq (2) and hence at a larger transition density. The detonation can be triggered locally when a region appears that is larger than the critical size and enters the distributed regime due to a strong *local* stretching. It is important to study the degree to which this intermittency effect increases the transition density, which is constrained by observations. Clearly the answer depends on the critical size, which we consider next.

The question of how large the isothermal region with well-mixed ash and fuel has to be for a detonation was studied by KOW (see also NW). In their model, the DDT occurs via the Zel'dovich mechanism (Zel'dovich et al. 1970) where the mixed region begins spontaneous ignition at the place with the minimum induction time, and the flame propagates with a phase speed equal to the inverse of the spatial gradient of the induction time, which is large for nearly isothermal and well-mixed regions and is not limited by the speed of sound. As the phase speed decreases below the Chapman-Jouget speed, a shock forms just ahead of the

flame front. Whether this shock can explode the whole star depends on the strength of the shock when entering the pure fuel, which is determined by the size of the isothermal region. If the isothermal region is small and the shock is weak, the flame front and shock separate with the flame front lagging behind the shock and the shock cannot make the whole star explode. The critical strength of the shock corresponds to a critical size of the isothermal region, over which the shock can be strengthened. Using 1D simulations, KOW and NW obtained the critical size, l_c , which depends on the density and the chemical composition. It is interesting to note that at early time when the density is large, the required size is much smaller than that at later times. We will show this has important consequences. The critical size is much smaller than what current numerical simulations can resolve, therefore the problem of the intermittent stretching at scale l_c cannot be addressed by simulations.

We need a local criterion to check whether a region of a given size l , in particular l_c , is in the distributed regime or not. For that purpose, we use a local average dissipation rate ϵ_l (see eq. 8 in §3 for a definition) in a region of size l to replace $\bar{\epsilon}$. Following the same argument that leads to equation (2), the criterion for a region of size l being in the distributed regime is

$$\epsilon_l > K^3 \epsilon_f \quad (3)$$

where we have used the refined similarity hypothesis by Kolmogorov (1962) (see eq. 9 in §3). Due to the random nature of turbulent flows, ϵ_l is stochastic, and a statistical approach is necessary. We therefore ask the question: what is the probability that any region of size l is in the distributed regime? This is given by the cumulative probability $P(\epsilon_l > K^3 \epsilon_f)$. To answer this question, we need the probability distribution $P(\epsilon_l)$ of ϵ_l . Fortunately, this distribution has been extensively studied in the intermittency models for turbulence, which we describe in §3. Although these models were originally proposed for homogeneous and isotropic turbulence, we will assume they apply to SNe Ia where the turbulence is stratified and may not arrive at isotropy even at very small scales. Once the distribution is specified, one can calculate the probability of finding that a region of given size l_c is in the distributed regime,

$$P(\epsilon_{l_c} > K^3 \epsilon_f) = \int_{K^3 \epsilon_f}^{\infty} P(\epsilon_{l_c}) d\epsilon_{l_c} \quad (4)$$

which depends on the density through l_c and ϵ_f . An immediate examination of eq (4) shows that, at larger density, the lower limit of the integral $K^3 \epsilon_f$ is larger because of the fast flame speed and the small flame thickness. This tends to decrease the probability. However, at larger density, l_c is smaller and the intermittency of turbulence tells us that the tail of the distribution $P(\epsilon_l)$ is broader for smaller l . This tends to counteract the decrease of the cumulative probability due to the larger lower integral limit at higher densities.

Furthermore, for smaller l_c , there are more regions of size l_c available in the star. This could make the transition occur significantly earlier with a transition density considerably larger than predicted by eq (2). We need to multiply the probability that a given region of size l_c is in the distributed regime by the number, N_{l_c} , of regions of size l_c available in order to calculate the number of regions that are both larger than l_c and in the distributed regime at any given density. We assume that the deflagration to detonation transition happens when

$$N_{l_c} \times P(\epsilon_{l_c} > K^3 \epsilon_f) = 1. \quad (5)$$

Since we are concerned with the flame being stretched into the distributed regime, only locations around the flame front are of interest when calculating N_{l_c} . Therefore, we only count regions in the vicinity of the flame front. N_{l_c} depends on the size, R_f , of the flame region and the flame geometry. A typical value for R_f is 10^8 cm (Khokhlov 1995), which could be smaller at an earlier time. We will set $R_f \simeq L$ in our calculations in order to decrease the number of parameters. Note that $R_f > L$ when the freezeout effect is considered and therefore the number N_{l_c} we use is a lower limit. If the flame region is a 2D spherical front, $N_{l_c} \sim 4\pi R_f^2/l_c^2$. If the flame structure is highly convoluted, it may have a fractal dimension larger than 2. In that case, N_{l_c} is larger. The upper limit for N_{l_c} is $\simeq 4\pi R_f^3/3l_c^3$, which applies if the flame geometry is close to 3D. Again, we take the lower limit $N_{l_c} = 4\pi R_f^2/l_c^2$, thus the transition density we will get is a lower limit.

As discussed in the Introduction, Lisewski et al. (2000) and Woosley (2007) find that entering the distributed regime is not sufficient for the DDT to occur and give criteria stronger than that used in KOW and NW. Lisewski et al. (2000) considered how turbulent transport affects the temperature and composition profile around a laminar flame. They assumed that, at any point, turbulence translates the temperature and composition by a distance l_t , over which turbulence can transport during a local induction time τ_i . The distance l_t is a function of position since τ_i depends on local temperature and composition. It is estimated by the length scale of a turbulent eddy with turnover time equal to τ_i , i.e., $l_t/\delta u(l_t) = \tau_i$. Using the Kolmogorov (1941) scaling, we get $l_t = \bar{\epsilon}^{1/2} \tau_i^{3/2}$. For given turbulence intensity, temperature and composition profiles around a laminar flame front can be calculated from the translation. Clearly, more efficient turbulent transport gives shallower temperature and composition profile, which is needed for detonation. By checking whether the resulting profiles, as initial conditions to solve the 1D hydrodynamic equations, can lead to a detonation, Lisewski et al. (2000), obtained a condition for the DDT on the turbulent intensity. They found that, for a successful detonation, the turbulent velocity has to be $\sim 10^8$ cm/s at the scale 10^6 cm. This condition is stronger than just entering the distributed

regime³. Since the expected turbulent velocity at scale 10^6 cm is $10^6 - 10^7$ cm/s, Lisewski et al. (2000) concluded that a DDT via the Zel’dovich mechanism in SNe Ia is unlikely. However, considering the spatial inhomogeneity of turbulent intensity, i.e., intermittency, it is possible for regions of size 10^6 cm with large enough turbulent velocity to arise.

The result of Lisewski et al. (2000) motivated Röpke (2007) to study the probability of finding a region of size 10^6 cm with a turbulent rms velocity of $\sim 10^8$ cm/s. Using data from 3D numerical simulations with a turbulent subgrid-scale method, Röpke (2007) analyzed the velocity fluctuations at the grid size (10^6 cm) and obtained a fat exponential tail for large velocity fluctuations that extends up to 10^8 cm/s. The large velocity fluctuations seem likely to be located at the trailing edge of a bubble-like feature (Röpke 2007). This confirms the intermittency in the turbulent combustion flow in SNe Ia; there exist grid cells where the turbulent intensity is much stronger than the average. From the probability of finding a grid cell with required turbulent intensity, Röpke concluded that the DDT triggered by a local cell with large velocity fluctuations is possible but probably rare. In our notations, the probability is given by $P(\epsilon_{10^6 \text{ cm}} > 10^{18} \text{ cm}^2/\text{s}^3)$ where $10^{18} \text{ cm}^2/\text{s}^3$ corresponds to the dissipation rate in a region of size 10^6 cm with a rms velocity of 10^8 cm/s. We will calculate this probability and consider the availability of such regions using two intermittency models given in §3 and compare with the results of Röpke (2007) in §4.

Woosley (2007) proposed a new criterion for the DDT based on a calculation of the distributed flame width using an eddy diffusivity approximation. Making an analogy to the estimate of the laminar flame thickness, Woosley (2007) obtained the distributed flame width λ from the equation $\lambda \simeq (D(\lambda)\tau_n)^{1/2}$ where $D(\lambda) = \delta u(\lambda)\lambda$ is the eddy diffusivity at scale λ and τ_n is the nuclear reaction timescale. Using the Kolmogorov (1941) scaling for $\delta u(\lambda)$, the distributed flame width is given by $\lambda = \bar{\epsilon}^{1/2}\tau_n^{3/2}$ (note that this formula for λ is similar to l_t in Lisewski et al. (2000)). Woosley (2007) assumed that the condition for detonation is that the minimum burning timescale in the distributed flame is smaller than the sound crossing time over the distributed flame width λ , or equivalently, $\lambda \gtrsim r_{sonic}^{min}$ where r_{sonic}^{min} is the sound crossing length over the minimum burning timescale in the distributed flame. The minimum sound crossing length is thus the critical size of the distributed flame width for detonation. The criterion $\lambda \gtrsim r_{sonic}^{min}$ is equivalent to $\bar{\epsilon} \gtrsim (r_{sonic}^{min})^2/\tau_n^3$. Noting that

³This condition can be converted into a form that can be directly compared with eq. 2. Roughly speaking, the physical condition for a detonation in this model is that l_t at the laminar flame front is larger than l_c , i.e., a shallow temperature gradient can be produced over a critical size around the flame front. Requiring $l_t > l_c$, we get the criterion $\bar{\epsilon} > (l_c/l_f)^2(\tau_n/\tau_i)^3\epsilon_f$. Considering that τ_i defined by Lisewski et al. (2000) is smaller than the nuclear timescale τ_n and that $l_c \gg l_f$, this condition is much stronger than the condition of eq. 2. Note that this condition is similar to the criterion of Woosley (2007) given below.

$\tau_n = l_f/S_l$, the condition can be written as $\bar{\epsilon} \gtrsim (r_{sonic}^{min}/l_f)^2 \epsilon_f$. Since r_{sonic}^{min} given in Table 4 of Woosley (2007) is close to l_c listed in Table 1 in this paper, we will use l_c instead of r_{sonic}^{min} for simplicity, i.e.,

$$\bar{\epsilon} \gtrsim (l_c/l_f)^2 \epsilon_f. \quad (6)$$

which is much stronger than eq (2) because l_c is much larger than the laminar flame thickness l_f . This condition can be used to determine the transition density ρ_{tr} by the same calculation process as in the case of the criterion eq (2). Note that, except a factor of $(\tau_n/\tau_i)^3$, this criterion is basically equivalent to that given in footnote (3) for the requirement by Lisewski et al. (2000). We find that, for the reasonable turbulence parameters listed in Table 2, the criterion results in a transition density below 10^7 g cm^{-3} and we cannot specify it due to the lack of data for l_c , S_l and l_f at densities below 10^7 g cm^{-3} .

In his estimate for ρ_{tr} , Woosley (2007) used $U = 10^8 \text{ cm/s}$ at scale $L = 10^6 \text{ cm}$ throughout the calculations, based on the result by Röpke (2007) on the possibility of the existence of regions of size 10^6 cm with a rms velocity of 10^8 cm/s . With these turbulence parameters he derived $\rho_{tr} = 10^7 \text{ g cm}^{-3}$. Clearly, in Woosley’s calculation, the intermittency effect implicitly contributes to the transition density obtained because, as discussed earlier, a turbulent rms velocity of 10^8 cm/s at 10^6 cm can only arise from intermittency.

The intermittency effect for the criterion of Woosley (2007) can be included more consistently in our formulation. Instead of considering a single special scale 10^6 cm , our model specifies intermittency over a continuous range of scales corresponding to critical sizes at different densities. Following the same steps that lead to eq (5), we incorporate the intermittency effect in the DDT model of Woosley (2007) and obtain a criterion,

$$N_{l_c} \times P(\epsilon_{l_c} > (l_c/l_f)^2 \epsilon_f) = 1. \quad (7)$$

which only differs from eq. 5 by the lower limit in the cumulative probability. We will discuss about this criterion in §4.

We point out that the eddy diffusivity method used by Woosley (2007) to approximate the combined action of the turbulent advection and the microscopic diffusivity is an oversimplification. This procedure implicitly assumes a smooth structure in the distributed flame and neglects the fluctuations of temperature and concentration, which may be important in determining the effective width of distributed flames.

3. Intermittency

Kolmogorov’s 1941 theory assumes that the energy transfer in the inertial range is equal to the average dissipation rate $\bar{\epsilon}$ in the flow and is the same throughout the inertial scales

down to the viscous scale where the kinetic energy is removed. This assumption, together with the similarity hypothesis, predicts that the statistics of the velocity difference (or the velocity fluctuations) at any inertial scale is completely determined by the average dissipation rate $\bar{\epsilon}$. However, fluctuations in the dissipation rate clearly exist as can be seen from the formula for the local viscous dissipation rate, $\epsilon(\mathbf{x}, t) = \frac{\nu}{2} \sum_{i,j} (\partial_i u_j + \partial_j u_i)^2$, which is a function of the fluctuating velocity field. The spatial fluctuations in ϵ are well-illustrated by the intense dissipation structures at small scales such as vortex tubes. This effect needs to be taken into account for a more accurate prediction of the scaling behavior of the velocity difference (Landau and Lifshitz 1944). The statistics of the velocity difference over a separation l depends on the distribution of the dissipation rate over regions of size l , which is defined as (e.g., Kolmogorov 1962),

$$\epsilon_l(\mathbf{x}, t) = \frac{3}{4\pi l^3} \int_{|\mathbf{x}'| < l} \epsilon(\mathbf{x} + \mathbf{x}', t) d\mathbf{x}'. \quad (8)$$

Clearly, the mean of ϵ_l is equal to $\bar{\epsilon}$ and thus is independent of l . This means that the *average* energy flux over all the inertial scales is constant. The ϵ_l distribution is essential to the intermittency models for turbulence. Note that this distribution is exactly what we need in our calculations for the transition of the turbulent combustion to the distributed regime by turbulent stretching and quenching described in §2, eqs (4) and (5).

Intermittency in turbulence is usually expressed in terms of the scaling behavior of the structure functions $\langle \delta u(l)^p \rangle \sim l^{\zeta_p}$ where $\delta u(l) = u(x+l) - u(x)$ is the (longitudinal) velocity difference and ζ_p is the scaling exponent for the p th-order structure function. Kolmogorov's 1941 theory predicts that the exponent ζ_p goes with p as $\zeta_p = p/3$. However experimental data (e.g., Anselmet et al. 1984) have shown departure from this linear relation and ζ_p increases significantly slower than $p/3$ at large p . This “anomalous” scaling is referred to as intermittency. The data indicate broader and broader tails for the distribution of $\delta u(l)$ at smaller and smaller scales, e.g., the kurtosis of the distribution, $\langle \delta u(l)^4 \rangle / \langle \delta u(l)^2 \rangle^2 \propto l^{\zeta_4 - 2\zeta_2}$, increases with decreasing l because $\zeta_4 < 2\zeta_2$. The distribution of $\delta u(l)$ is fatter for smaller l . The anomalous scaling is fundamentally caused by the fluctuations in the dissipation rate ϵ_l . Applying the refined similarity argument for homogeneous and isotropic turbulence (Kolmogorov 1962), the velocity difference over a separation l can be related to the dissipation rate ϵ_l ,

$$\delta u(l) \sim \epsilon_l^{1/3} l^{1/3}. \quad (9)$$

(note that the Kolmogorov's 1941 theory uses $\bar{\epsilon}$.) The structure functions are then given by,

$$\langle \delta u(l)^p \rangle \propto \langle \epsilon_l^{p/3} \rangle l^{p/3}. \quad (10)$$

Clearly, the departure from the linear scaling for the velocity difference comes from the statistics of the dissipation rate. Assuming $\langle \epsilon_l^p \rangle \propto l^{\tau_p}$ (e.g., She and Leveque 1994), we have,

$$\zeta_p = p/3 + \tau_{p/3}. \quad (11)$$

Developing a physical model for τ_p that satisfies the experimental result for $\zeta(p)$ has been the main task of intermittency theories. Although we are mainly concerned with the distribution of ϵ_l , discussions of the structure functions are necessary because they are directly measurable in experiments and give important information and constraints on the ϵ_l distribution. We will use two intermittency models in our calculations: the log-normal model (Oboukhov 1962 and Kolmogorov 1962) and the log- Poisson model by She and Leveque (1994).

3.1. The log-normal model

Oboukhov (1962) and Kolmogorov (1962) developed the first intermittency model. In this model, the distribution of ϵ_l is assumed to be log-normal (Kolmogorov 1962). A justification for this “natural” distribution for ϵ_l was given by Yaglom (1966). Imagine the cascade progress as successive eddy fragmentations from the integral scale L to the dissipation scale η . The statistics of the energy flux at an inertial scale l (or equivalently the dissipation rate ϵ_l) depends on the fragmentations before the scale is reached. The total number N of steps that lead to the scale l is proportional to $N \sim \ln(L/l)$. Defining $\chi_i = \epsilon_i/\epsilon_{i-1}$ as the ratio of the energy transfer rates at two successive fragmentation steps, the energy flux at the scale l can be expressed in the ratios (see e.g., Monin & Yaglom 1975),

$$\epsilon_l = \epsilon_L \chi_1 \chi_2 \dots \chi_N \quad (12)$$

where ϵ_L is the dissipation rate at the integral scale (or the transfer flux at the largest scale), $\epsilon_L \simeq \bar{\epsilon}$. Due to the randomness in the fragmentation process, χ_i ’s are stochastic variables. Assuming a self- similar fragmentation process, the distributions of χ_i ’s are similar and $\ln(\epsilon_l/\epsilon_L) = \sum_{i=1}^N \ln(\chi_i)$ is expected to be Gaussian from the central limit theorem,

$$p(\epsilon_l)d\epsilon_l = \frac{1}{\sqrt{2\pi\sigma_l^2}} \exp\left(-\frac{(\ln(\epsilon_l/\bar{\epsilon}) + \sigma_l^2/2)^2}{2\sigma_l^2}\right) d\ln(\epsilon_l/\bar{\epsilon}) \quad (13)$$

where the variance σ_l^2 is proportional to the number of steps $\sigma_l^2 = \mu \ln(L/l)$ with μ being a parameter to be determined by experimental data and the $\sigma_l^2/2$ term in the numerator in the exponential is to guarantee the mean $\langle \epsilon_l \rangle$ is equal to the overall average dissipation rate $\bar{\epsilon}$. This distribution will be used later to calculate the probability (eq. 4) for a region of a given size being in the distributed regime.

The scaling behavior of ϵ_l can be derived by integrating equation (13),

$$\langle \epsilon_l^p \rangle \propto l^{-\frac{1}{2}\mu p(p-1)} \quad (14)$$

which gives $\tau_p = -\frac{1}{2}\mu p(p-1)$. From eq (11), we have

$$\zeta_p = p/3 - \frac{1}{18}\mu p(p-3) \quad (15)$$

Therefore $\mu = 2 - \zeta_6$, which can be obtained from the results of experiments and simulations. It has been found that $\mu \simeq 0.2$ (Frisch 1995, Biskamp 2000). The relation (15) agrees with experiments quite well at small p but starts to exhibit deviation at $p \gtrsim 10$ and gives an unrealistic maximum and turnover at $p > 16$, violating the requirement that the $\zeta(p)$ - p curve must be monotonic and concave (Frisch 1995). Simulations by Wang et al. (1996) suggest that this disagreement corresponds to the departure of the distribution for $\ln(\epsilon_l/\bar{\epsilon})$ from normal at scales close to the dissipation scale. They find that, at these scales, the distribution of $\ln(\epsilon_l/\bar{\epsilon})$ shows a negative skewness, meaning that the log-normal distribution overestimates the probability in the very high ϵ_l tail. However the distribution of ϵ_l agrees with log-normal very well in the inertial range away from the dissipation scale and the agreement is better and better for larger and larger scales (Wang et al. 1996). Fortunately the critical scale we are concerned with is well within the inertial range (see §4) and according to Fig. 6 in Wang et al. (1996) the log normal fit is very good at least up to the $4 - \sigma$ tail. They also show that the fit gets better as the Reynolds number increases. More recent simulations by Yeung et al. (2006) with resolutions up to 2048^3 obtained similar results. The log normal distribution gives a very good fit to $4 - \sigma$ and only deviates by a factor of 2 at the $5 - \sigma$ tail. They also find that the negative skewness gets closer to zero with increasing Reynolds number. The Reynolds number in SNe Ia is $Re \sim 10^{14}$ for typical velocity scale 10^7 cm/s, length scale 10^7 cm and viscosity $1 \text{ cm}^2/\text{s}$. This is much larger than in all the current simulations. Therefore one may expect that the log-normal distribution probably applies even further out on the tail for the inertial scales of the turbulence in SNe Ia. However, the departure of the predicted $\zeta(p) - p$ curve from the experiments (with high Re) at $p \gtrsim 10$ suggests that, even at huge Reynolds number, the log normal fit eventually breaks down at some large ϵ_l in the tail. Therefore we need to be careful when using the log-normal model. We will give more discussion on this point in the calculations given in §4.

Another issue is that the distribution of ϵ_l has a physical cutoff in a realistic system due to the finite viscosity. Since the intermittency is stronger at smaller scales, the cutoff in the distribution of ϵ_l is probably larger for smaller l and obtains a maximum at the dissipation scale, η . For Kolmogorov scaling, the cutoff in the distribution of ϵ_η is given by $\bar{\epsilon} Re^{1/2}$. Since $Re \sim 10^{14}$ in SNe Ia, this maximum dissipation rate is far beyond that required to break flames at density $\lesssim 10^8 \text{ g/cm}^3$. Therefore ignoring this maximum cutoff does not

affect our result. However, the largest available dissipation rate at an inertial scale l is probably smaller than the cutoff in the distribution of ϵ_η and thus may affect the calculation for the cumulative probability defined in eq (4) if the cutoff in the distribution $P(\epsilon_{l_c})$ is close to or even smaller than $K^3\epsilon_f$. Since the log-normal model does not address the cutoff in the distribution of ϵ_l , we will neglect this potential effect in this model.

On the other hand, the log-Poisson model we consider in the next section gives a maximum dissipation rate at each inertial scale, corresponding to the strongest dissipative structures at that scale. In that model, a nonzero cumulative probability in eq (5) requires the lower limit $K^3\epsilon_f$ in eq (4) be smaller than the maximum.

3.2. The log-Poisson model

A major success in the intermittency theory is the model by She and Leveque (1994). In this model, She and Leveque studied the hierarchy of dissipation intensity in structures of size l and, by invoking an unknown “hidden symmetry,” they related the characteristic dissipation rates in structures of different intensity levels to the strongest dissipative structures. This relation gives a prediction of τ_p as a function of p , which only depends on the properties of the most intermittent structures. Assuming that the dissipation rate in regions of size l containing the most intense structures exhibits a scaling $\propto l^{-2/3}$ (see explanation in Appendix A) and the most intermittent structures are filamentary, corresponding to a codimension of 2, She and Leveque obtained a $\zeta(p) - p$ relation, which is in excellent agreement with experimental data. The “hidden symmetry” has been immediately interpreted as a log-Poisson process (Dubrulle 1994, She and Waymire 1995) in a multiplicative cascade model. In this section, we adopt the log-Poisson version of the She-Leveque model. The original presentation by She and Leveque (1994) is given in Appendix A.

In a multiplicative model, the dissipative rates at two scales l_2 and l_1 ($l_1 > l_2$) are related by a multiplicative factor $W_{l_1l_2}$,

$$\epsilon_{l_2} = W_{l_2l_1} \epsilon_{l_1} \quad (16)$$

The average $\langle W_{l_1l_2} \rangle$ is equal to unity since $\langle \epsilon_{l_1} \rangle = \langle \epsilon_{l_2} \rangle = \bar{\epsilon}$. She and Waymire (1995) speculated that $W_{l_2l_1}$ consists of two events. First is the amplification of the dissipation rate in the cascade, which tends to produce singular structures with $\epsilon_{l_2} \propto (l_1/l_2)^\gamma$ approaching infinity as l_2 goes to 0. The meaning of γ is discussed below. To ensure $\langle W_{l_1l_2} \rangle = 1$, a second event is required to reduce $W_{l_1l_2}$. She and Waymire (1995) called this event the modulation-defects since it modulates the singular structures. The defects were assumed to be a discrete

Poisson process. Each of the defects decreases $W_{l_1 l_2}$ by a factor of β , thus

$$W_{l_2 l_1} = (l_1/l_2)^\gamma \beta^n \quad (17)$$

if there are n defect events in the cascade. The number n of the events that occur in the cascade from the scale l_1 to l_2 obeys a Poisson distribution,

$$P(n) = \exp(-\lambda_{l_1 l_2}) \frac{\lambda_{l_1 l_2}^n}{n!} \quad (18)$$

where $\lambda_{l_1 l_2}$ is the mean number of the defect events in the cascade, which is expected to be proportional to the total number of the cascade steps, i.e., $\lambda_{l_1 l_2} \propto \ln(l_1/l_2)$. In fact, $\lambda_{l_1 l_2}$ can be obtained by taking the average of eq. (17) and requiring $\langle W_{l_1 l_2} \rangle = 1$. Using the identity $\sum_{n=0}^{\infty} \frac{\alpha^n}{n!} = \exp(\alpha)$, we get $\langle \beta^n \rangle = \exp((\beta - 1)\lambda_{l_1 l_2})$ for the Poisson distribution eq (18), therefore,

$$\lambda_{l_1 l_2} = \frac{\gamma \ln(l_1/l_2)}{1 - \beta} \quad (19)$$

In this model, there is a largest dissipation rate at any scale. Clearly the largest dissipation rate is achieved if there is no defect, i.e., $n = 0$, in a cascade from the integral scale L to the scale l of interest, thus the largest dissipation rate is equal to $\epsilon_L (L/l)^\gamma$. This largest dissipation rate corresponds to $\epsilon_l^{(\infty)}$ in Appendix A. Similarly $n = 1$ gives the second strongest dissipative rate at a given scale, and so on.

From eqs (16) and (17), we have,

$$\ln(\epsilon_{l_2}/\bar{\epsilon}) = \ln(\epsilon_{l_1}/\bar{\epsilon}) + \gamma \ln(l_1/l_2) + n \ln(\beta) \quad (20)$$

thus, using the Poisson distribution for n , the distribution for the dissipation rate at l_2 can be derived from that at any scale l_1 larger than l_2 . In particular, we consider deriving the distribution of ϵ_l at any scale l from the integral scale L . The distribution function of ϵ_L at the integral scale depends on how the energy is injected in the flow, thus is not universal and may vary from flow to flow. Therefore the function form cannot be specified. However, there is a strong constraint for its width. Since $\epsilon_L \simeq \bar{\epsilon}$, the distribution of $\ln(\epsilon_L/\bar{\epsilon})$ is expected to be very narrow around $\ln(\epsilon_L/\bar{\epsilon}) \simeq 0$ and hence to be approximately a delta function. We denote the distribution of $\ln(\epsilon_L/\bar{\epsilon})$ as $P_L(\ln(\epsilon_L/\bar{\epsilon}))$. It then follows from eqs (18) and (20) that

$$P(\epsilon_l) d\epsilon_l = \sum_{n=0}^{\infty} \exp(-\lambda) \frac{\lambda^n}{n!} P_L(\ln(\epsilon_l/\bar{\epsilon}) - \gamma \ln(L/l) - n \ln(\beta)) d\ln(\epsilon_l/\bar{\epsilon}) \quad (21)$$

where $\lambda = \lambda_{Ll} = \gamma \ln(L/l)/(1 - \beta)$. Each term in eq (21) represents the contribution from dissipation structures of different levels, e.g., the $n = 0$ term corresponds to the most intensive structures of size l .

To compare the model with experiments and obtain the parameters, we calculate the moments $\langle \epsilon_l \rangle$ from the distribution eq. (21),

$$\begin{aligned}
\langle \epsilon_l^p \rangle &\propto \sum_{n=0}^{\infty} \int \exp(-\lambda) \frac{\lambda^n}{n!} \exp(px) P_L(x - \gamma \ln(L/l) - n \ln(\beta)) dx \\
&= (l/L)^{-\gamma p} \exp(-\lambda) \sum_{n=0}^{\infty} \frac{(\beta^p \lambda)^n}{n!} \int \exp(x') P_L(x') dx' \\
&= B_p \exp(-\lambda(1 - \beta^p)) (l/L)^{-\gamma p} \\
&= B_p (l/L)^{-\gamma p + \gamma(1 - \beta^p)/(1 - \beta)}
\end{aligned} \tag{22}$$

where we used a variable change $x' = x - \gamma \ln(L/l) - n \ln(\beta)$ in the second step and the identity $\sum_{n=0}^{\infty} \frac{\alpha^n}{n!} = \exp(\alpha)$ in the third step. The coefficients $B_p = \int \exp(px') P_L(x') dx'$; $B_0 = B_1 = 1$ from the normalization of P_L and the requirement that $\langle \epsilon_L \rangle = \bar{\epsilon}$ respectively.

The result eq (22) gives $\tau_p = -\gamma p + \gamma(1 - \beta^p)/(1 - \beta)$, which is the same as (A8) in Appendix A, meaning that the “hidden symmetry” described in the appendix is equivalent to a log-Poisson process. The parameters γ and β introduced here are identical to those described in the appendix, and thus have the physical meanings explained there, i.e., γ can be interpreted as the exponent of the dissipation rate scaling in regions containing the most intermittent structures and β is related to the codimension C of the strongest dissipation structures, $\gamma/(1 - \beta) = C$ (see Appendix A for details). As discussed in the appendix, She and Leveque argued that $\gamma = 2/3$ and $\beta = 2/3$ for $C = 2$ corresponding to filamentary dissipation structures in incompressible turbulence. This results in ζ_p as a function of p that agrees with the experiments with an accuracy of 1%, implying eq (21) provides a good distribution for ϵ_l . The She-Leveque formulation has been extended to supersonic turbulence (Boldyrev et al. 2002) and MHD turbulence (Muller & Biskamp 2000) where the dissipation structures are dissipation sheets and the current sheets, respectively. For these 2 dimensional dissipation structures, the codimension $C = 1$ and $\beta = 1/3$. In next section we use the log-Poisson distribution (eq 21) in our calculations for the cumulative probability in eq 4. We will take $\gamma = 2/3$ and consider both filaments ($\beta = 2/3$) and sheets ($\beta = 1/3$) as the most intermittent dissipation structures.

4. Results

4.1. The log-normal model

We are ready to calculate the probability $P(\epsilon_{l_c} > K^3 \epsilon_f)$ using the distributions $P(\epsilon_l)$ given in §3. The calculation is straightforward for the log-normal distribution eq. (13),

$$\begin{aligned}
 P(\epsilon_{l_c} > K^3 \epsilon_f) &= \int_{\ln(K^3 \epsilon_f / \bar{\epsilon})}^{\infty} \frac{1}{\sqrt{2\pi\sigma_{l_c}^2}} \exp\left(-\frac{(x + \sigma_{l_c}^2/2)^2}{2\sigma_{l_c}^2}\right) dx \\
 &= \int_{\frac{\ln(K^3 \epsilon_f / \bar{\epsilon}) + \sigma_{l_c}^2/2}{\sigma_{l_c}}}^{\infty} \frac{1}{\sqrt{2\pi}} \exp(-x^2/2) dx \\
 &= \frac{1}{2} \operatorname{erfc}\left(\frac{\ln(K^3 \epsilon_f / \bar{\epsilon})}{\sqrt{2}\sigma_{l_c}} + \frac{\sigma_{l_c}}{2\sqrt{2}}\right)
 \end{aligned} \tag{23}$$

where $\bar{\epsilon} \simeq U^3/L$, $\sigma_{l_c}^2 = \mu \ln(L/l_c) \simeq 0.2 \ln(L/l_c)$ and $\operatorname{erfc}(x)$ is the complementary error function. Using ϵ_f and l_c given in Table 1, we calculated the probability as a function of the density assuming different values for the characteristic velocity (U) and length (L) scales. For example, if $U = 100$ km/s and $L = 100$ km, the probabilities are 0.5×10^{-10} , 0.042, 0.9 and 1 at $\rho = 10^8$, 5×10^7 , 3×10^7 and 10^7 g/cm³ respectively if $K = 1$. It is interesting to note that, at a density 3×10^7 g/cm³, 10% of the local regions of the critical size are still in the flamelet regime, although generally the flame has reached the distributed regime according to the mean criterion eq (2). With $R_f \simeq L = 100$ km, at the 4 densities above from high to low, the corresponding numbers $N_{l_c} = 4\pi R_f^2/l_c^2$ of regions of the critical size that cover the flame front are 3×10^{10} , 7×10^8 , 5×10^7 and 3×10^4 . Multiplying $P(\epsilon_{l_c} > K^3 \epsilon_f)$ with N_{l_c} (eq. 5), we see that there is already one region of size l_c in the distributed regime at a density 10^8 g/cm³. Recalling that, according to the criterion eq (2), the DDT does not occur until the density decreases to 4×10^7 g/cm³, we find that in this case the intermittency effect may increase the transition density by more than a factor of 2.

We point out the cumulative probability calculated from eq (23) at density 10^8 g/cm³ in the example above comes from a little beyond the $6 - \sigma$ tail of the distribution for $\ln(\epsilon_l/\bar{\epsilon})$ and we need to check whether the log-normal distribution there is a good approximation. As discussed in §3.1, numerical simulations have shown that for a scale l in the inertial range, the distribution of ϵ_l is well approximated by log-normal up to the $5 - \sigma$ tail (Yeung et al. 2006). Assuming the Kolmogorov scaling, the dissipation scale in Type Ia SNe is $\eta = LRe^{-3/4} \simeq 10^{-3}$ cm for $L \sim 10^7$ cm and the Reynolds number $Re \simeq 10^{14}$. The critical scale $l_c \sim 10^2 - 10^4$ cm of interest here is well between the integral scale and the dissipation scale, thus we expect that the distribution for ϵ_{l_c} is close to log-normal at least up to the $\sim 5 - \sigma$ tail. The question is then whether the good fit extends further. Wang et al. (1996) and Yeung et al. (2006) found that the log-normal fit is better for larger Reynolds

number, thus it is expected that the log-normal approximation probably applies to higher on the tail than $5 - \sigma$. As argued in §3.1, the log normal approximation eventually fails somewhere in the extreme tail even at high Reynolds number. To know exactly how far the log-normal fit extends, numerical simulations with much higher resolution are needed. We have to be careful about the validity of the log-normal approximation in the far tail because it overestimates the probability distribution for ϵ_l once it breaks down and in that case eq (23) overestimates $P(\epsilon_{l_c} > K^3 \epsilon_f)$.

Due to the complication of the validity of the log-normal distribution at the far tail, we consider two extreme cases and give the upper and lower limits for the transition density. First, we ignore the departure from log-normal and evaluate the density at which $N_{l_c} \times P(\epsilon_{l_c} > K^3 \epsilon_f) = 1$ using eq (23) for $P(\epsilon_{l_c} > K^3 \epsilon_f)$ and Table 1 for ϵ_f and l_c with different parameters U and L . We will denote this density as ρ_{LN} with the subscript LN standing for log-normal. Interpolation was used to obtain ϵ_f and l_c not tabulated in Table 1. If the distribution of ϵ_l is exactly log-normal as given by eq (13), then ρ_{LN} is the predicted transition density for the DDT with the intermittency taken into account. On the other hand, if the log-normal distribution overestimates the probability at the high tail, eq (23) overestimates the cumulative probability and ρ_{LN} is the upper limit for ρ_{tr} . We give ρ_{LN} for different parameters U and L in the second line of Table 2.

In the other extreme, we assume that the log-normal distribution fails to fit the distribution of ϵ_l beyond the $5 - \sigma$ tail. This gives a lower limit for the transition density since numerical simulations have shown that the log-normal fit applies at least to $5 - \sigma$. In this case, we keep track of the integral limit in the second line of eq (23) at ρ_{LN} , which tells us which part of the tail of the distribution gives the main contribution to $P(\epsilon_{l_c} > K^3 \epsilon_f)$ at that density. If the integral limit is smaller than 5, the contribution to the cumulative probability is from within $5 - \sigma$ and vice versa. We calculate the density at which the integral limit is equal to 5 and denote this density as $\rho_{5\sigma}$. Since the integral limit is a decreasing function of the density, if $\rho_{LN} < \rho_{5\sigma}$, the contribution to $P(\epsilon_{l_c} > K^3 \epsilon_f)$ at density ρ_{LN} is from within $5 - \sigma$. In this case, the cumulative probability calculated from eq (23) is valid and ρ_{LN} is a good estimate for the transition density. Otherwise if $\rho_{LN} > \rho_{5\sigma}$, the contribution to the probability is from beyond the $5 - \sigma$ tail, eq (23) overestimates it and thus ρ_{LN} overestimates the transition density ρ_{tr} . In this case, $\rho_{5\sigma}$ gives a lower limit for the transition density because at $\rho_{5\sigma}$, we have $N_{l_c} \times P(\epsilon_{l_c} > K^3 \epsilon_f) \gg 1$ using eq. (23) which applies for $\rho \leq \rho_{5\sigma}$. Therefore if the log-normal fit fails just beyond $5 - \sigma$, we have a lower limit for the transition density, $\min(\rho_{LN}, \rho_{5\sigma})$. We give this lower limit in the 3rd line of Table 2.

Similar calculations can be done for the $K = 8$ case. The results of the upper and lower limits for the transition density in the $K = 8$ case are given in parenthesis in Table

2. Comparing with predictions from the mean criterion eq (2) (the first line in Table 2), the log-normal model predicts that the intermittency effect increases the transition density by a factor of 2-3 for all the cases we list in Table 2.

We evaluate the probability of the existence of a region of size 10^6 cm with a rms velocity 10^8 cm/s, required for the DDT by Lisewski et al. (2000), and compare with the numerical results of Röpke (2007). Using the log-normal distribution for ϵ_l , we find that the requirement $P(\epsilon_{10^6 \text{ cm}} > 10^{18} \text{ cm}^2/\text{s}^3)$ requires conditions from the extreme tail of the distribution. The likelihood is completely negligible ($\sim 10^{-40}$) if the velocity U at the integral length scale is less than $\sim 10^7$ cm/s. Only if U is larger than 5×10^7 cm/s is the probability appreciably larger so that the required region might be available. For example, if $U = 5 \times 10^7$ cm/s at $L = 10^7$ cm, the probability is $\sim 10^{-11}$. This is still too small to guarantee the existence of a region as required by Lisewski et al. (2000). The number of available candidate regions of size 10^6 cm around the flame front is probably smaller than $10^5 - 10^7$, assuming the flame front radius is $\sim 10^8 - 10^9$ cm. This result agrees with the conclusion of Röpke (2007) that the existence of a region as required by Lisewski et al. (2000) is rare. To ensure such a region, the velocity at the integral scale has to be larger than 10^8 cm/s, which is probably impossible as discussed in §2.

We also carry out a calculation for ρ_{tr} based on the criterion of Woosley (2007) taking into account the effect of intermittency. Using the log-normal distribution to calculate the cumulative probability in eq. (7), we find that no regions of critical size that meet Woosley’s criterion appear at density above 10^7 g/cm³. We cannot give an exact predicted transition density for this model because we do not have data at densities below 10^7 g/cm³ for relevant quantities listed in Table 1. Note that Woosley (2007) obtained a transition density around 10^7 g/cm³ under the assumption that a region of size 10^6 cm with rms velocity of 10^8 cm/s is available. From our estimate above and the result in Röpke (2007), the probability that such a region exists is small, therefore it is appropriate to take the transition density predicted in Woosley (2007) as an upper limit for his DDT criterion.

4.2. The log-Poisson model

We next consider the log-Poisson model. Using the distribution eq (21), we have,

$$\begin{aligned}
 P(\epsilon_{l_c} > K^3 \epsilon_f) &= \exp(-\lambda_c) \sum_{n=0}^{\infty} \frac{\lambda_c^n}{n!} \int_{\ln(K^3 \epsilon_f / \bar{\epsilon})}^{\infty} P_L(x - \gamma \ln(L/l_c) - n \ln(\beta)) dx \\
 &= \exp(-\lambda_c) \sum_{n=0}^{\infty} \frac{\lambda_c^n}{n!} \int_{\ln(K^3 \epsilon_f / \bar{\epsilon}) - \gamma \ln(L/l_c) - n \ln(\beta)}^{\infty} P_L(x) dx \\
 &= \exp(-\lambda_c) \sum_{n=0}^{\infty} \frac{\lambda_c^n}{n!} F_n
 \end{aligned} \tag{24}$$

where $\lambda_c = \gamma \ln(L/l_c)/(1 - \beta)$ and the integrals in the second line are denoted as F_n for convenience. Note that the integral lower limit $\ln(K^3 \epsilon_f / \bar{\epsilon}) - \gamma \ln(L/l_c) - n \ln(\beta)$ increases with n because $\beta < 1$, therefore F_n is a decreasing function of n . An exact calculation for the cumulative probability is impossible because of the unspecified function P_L . We will neglect all the $n \geq 1$ terms and only keep the $n = 0$ term in our calculation, i.e., we only include the contribution of the most intensive structures at scale l_c . Obviously, this approximation gives a lower limit for the probability and the transition density we obtain will also be a lower limit. We will show that the criterion for DDT obtained from this approximation is exact if P_L is a delta function.

The contribution from $n = 0$ is $\exp(-\lambda_c) = (\frac{l_c}{L})^{\gamma/(1-\beta)} F_0$. For $\gamma = 2/3$ and $\beta = 2/3$, it is equal to $(l_c/L)^2 F_0$ and $N_{l_c} \times P(\epsilon_{l_c} > K^3 \epsilon_f) \geq 4\pi(R_f/l_c)^2 (l_c/L)^2 F_0 = 4\pi(R_f/L)^2 F_0$. Since the size of the flame region $R_f \geq L$, it means that the number of regions which are larger than the critical size and in the distributed regime is $\simeq 4\pi F_0$. Since the distribution $P_L(x)$ is probably strongly concentrated at $x = 0$, the sufficient and almost necessary condition for $F_0 \simeq 1$ is that the integral limit $\ln(K^3 \epsilon_f / \bar{\epsilon}) - \gamma \ln(L/l_c) \leq 0$, or equivalently,

$$\bar{\epsilon} > (l_c/L)^{2/3} K^3 \epsilon_f \tag{25}$$

which is a convenient criterion for the DDT in the log-Poisson model. Note this criterion is much weaker than the mean criterion eq (2). Once the condition is satisfied, at least one region of critical size that covers the flame enters the distributed regime due to the most intense stretching strength available at scale l_c .

As mentioned in §3.2, if the dissipation structures are 2-dimensional, $\beta = 1/3$. In that case, the contribution from the $n = 0$ term is $(l_c/L) F_0$ and $N_{l_c} \times P(\epsilon_{l_c} > K^3 \epsilon_f) \geq 4\pi(R_f/l_c)(R_f/L) F_0$, which is much larger than 1 if $F_0 \gtrsim 1$. Therefore, the criterion eq (25) is a sufficient condition for the case with sheet-like dissipation structures such as in MHD turbulence or highly compressible turbulence.

We have neglected the $n > 1$ terms in eq. (24), the contribution of which depends on how rapidly $P_L(x)$ decreases with $x > 0$. We consider the extreme example where P_L is a

delta function. In this case, before the condition eq (25) is met, $F_n = 0$ for any n thus the cumulative probability is zero. When the condition is just satisfied as the density decreases, only the $n = 0$ term contributes and all the $n > 1$ terms are still zero, i.e., the most intensive ($n = 0$) structures at l_c can stretch a local flame into the distributed regime while all the less intensive structures ($n \geq 1$) still cannot. From the calculation above, we see that in this case once the $n = 0$ term contributes, at least one region around the flame front experiences the largest stretching rate and enters the distributed regime. Therefore, if P_L is a delta function, eq (25) is both the necessary and the sufficient condition. This is true for both $\beta = 2/3$ and $\beta = 1/3$. If P_L is not a delta function, the tail of P_L gives rise to the possibility that the distributed regime can emerge in a local region of critical size before the condition eq (25) is met. This could lead to an even weaker condition than eq (25). Since we expect that ϵ_L can only vary within a factor of a few, the condition can be weaker only by a factor of a few. Because the r.h.s of the condition (23), especially ϵ_f , depends on the density very sensitively, this would not increase the predicted ρ_{tr} considerably.

The condition eq (25) can be easily applied to calculate the transition density using ϵ_f and l_c in Table 1. For example, we get $\rho_{tr} = 8.7 \times 10^7$ g/cm³ for $U = 100$ km/s and $L = 100$ km if $K = 1$. This result is consistent with that from the log-normal model and is also about a factor of 2 larger than the prediction by the mean criterion eq (2).

The transition density predicted by the log-Poisson model with different parameters for turbulence is given in the 4th line of Table 2. Again the numbers in parenthesis are for $K = 8$. The results are consistent with those from the log-normal model and are at least 2 – 3 times larger than from the criterion eq (2).

Again we consider the possibility that there exists a region of size 10^6 cm with rms turbulent velocity of 10^8 cm/s required for DDT by Lisewski et al. (2000). The probability $P(\epsilon_{10^6 \text{ cm}} > 10^{18} \text{ cm}^2/\text{s}^3)$ depends on P_L , the probability distribution of the dissipation rate at the integral scale L . Since P_L is probably not universal and is flow-dependent, the log-Poisson model cannot give an exact estimate for the probability. Here we assume P_L is a delta function and see under what condition it is possible to find a required region. We find that the necessary and sufficient condition to have such a region is that $U > 10^{22/3}(L/\text{cm})^{1/9}$ cm/s. For $L \simeq 10^7$ cm, U has to be larger than 10^8 cm/s. This can be understood from the fact that, in the log-Poisson model, the available kinetic energy in the most intermittent structures for dissipation is assumed to be the kinetic energy at the integral scale (see Appendix A). Since $U > 10^8$ cm/s is probably not achievable, it is rare that a region as required by Lisewski et al. (2000) exists, again in agreement with Röpke (2007).

Using the same calculation that leads to eq (25), we obtain a criterion for the DDT

model by Woosley (2007) accounting for the intermittency effect,

$$\bar{\epsilon} > (l_c/L)^{2/3}(l_c/l_f)^2\epsilon_f. \quad (26)$$

This is weaker than the corresponding mean criterion eq (6) by a factor of $(l_c/L)^{2/3}$, meaning that intermittency increases the transition density. In comparison with eq (25), the condition is stronger and thus gives a smaller transition density than that for the KOW and NW model with intermittency included. At smaller density, the critical length l_c is larger and the factor $(l_c/L)^{2/3}$, representing the intermittency effect, is closer to unity. This implies that intermittency gives a weaker effect on the transition density for the Woosley (2007) criterion than for that by KOW and NW. Using Table 1, we again find that the condition eq (26) is not satisfied at densities above 10^7 g/cm³ for the five cases listed in Table 2, i.e., the predicted transition density is still below 10^7 g/cm³ after including intermittency (see discussion in §4.1).

In summary, intermittency can considerably enhance the onset of the distributed flame regime and hence increase the transition density in the DDT model of KOW and NW. Both the intermittency models we consider here predict a transition density 2–3 times larger than from the criterion using the mean dissipation rate. This factor of 2–3 brings the transition density to be in disagreement with the observational constraints for turbulent velocity larger than $U = 10^6$ cm/s in the case $K = 1$. We discuss the implications of this result in the next section. We also find that existence of regions of size 10^6 cm with velocity 10^8 cm/s is rare, in agreement with the numerical result of Röpke (2007). The strong DDT criterion given by Woosley (2007) gives a transition density below $\rho_{tr} = 10^7$ g/cm³ even when intermittency is included. We expect that the intermittency effect is weaker for stronger DDT criteria.

5. Conclusion and Discussion

We have studied the effect of intermittency on the transition from the flamelet regime to the distributed regime in Type Ia SNe, and hence on the transition density for the DDT model by KOW and NW. In their model, the detonation occurs via the Zel’dovich mechanism that requires a nearly isothermal region larger than a critical size to drive a sufficiently strong supersonic shock. KOW and NW assumed that the almost isothermal mixture of fuel and ash can be produced once turbulence is strong enough to get the flame into the distributed regime. The DDT is assumed by KOW and NW to occur when the *average* flow gets into the distributed regime. We argue that the sufficient condition for the DDT is that there is *one* region that is larger than the critical size and in the distributed regime.

The intermittency in turbulence, as a result of the spatial inhomogeneity of the dissipation rate, gives rise to regions with strong local turbulent strength that can force the flame into the distributed regime earlier than elsewhere. Therefore the transition from the flamelet regime to the distributed regime is not spatially smooth, but intermittent. At early time when the density in the white dwarf is large, the flame has a large speed and a small width and thus resists being efficiently stretched and broken by the turbulence. At the same time, the critical size is very small. This has two effects that tend to make an early DDT likely. First, the intermittency of turbulence tells us that the probability of finding extremely strong stretching within a smaller critical size is larger. Second, there are more regions of smaller sizes available. Therefore it is possible that the DDT is triggered at a small “spot” when the density is larger than needed for the average flow to enter the distributed regime. As we pointed out in the Introduction, the critical size as a function of the density plays an important role in determining the transition density for the DDT in our calculations.

We used two analytical intermittency models to statistically investigate when the first region appears which is both larger than the critical size and in the distributed region. This is assumed to be the time when the DDT occurs by KOW and NW. We found that, for various parameters for the intensity and length scales, DDT occurs at a transition density at least 2 – 3 times larger than the density at which the average flow enters the distributed regime. The transition density has been determined empirically by invoking it as a free parameter in spherically-symmetric models and then computing models that best match the observed multicolor light curve shapes and magnitudes (Höflich & Khokhlov 1996). Recognizing that the spherical models are oversimplified, they do give some guidance to the empirical constraints on the density at which DDT occurs. Höflich (1995) used this procedure to fit observations of the Branch core normal SN 1994D and preferred a value of the transition density of $2 \times 10^7 \text{ gm cm}^{-3}$. Höflich, Khokhlov & Wheeler (1995) explored a range of transition densities in the context of pulsating delayed detonation models and favored densities in the range $0.8 - 2.2 \times 10^7 \text{ gm cm}^{-3}$. Dominguez, Höflich & Straniero (2001) adopted $2.3 \times 10^7 \text{ gm cm}^{-3}$. Allowing for an uncertainty of a factor of 2, the predicted transition densities by the mean criterion are consistent with $2 \times 10^7 \text{ gm cm}^{-3}$ as favored by the observations in all the cases except that with $\bar{\epsilon} = 10^{16} \text{ cm}^2/\text{s}^3$ and $K = 1$ (Table 2). With the intermittency effects we have examined here, the transition density would be a factor of 2 – 3 higher. If $K = 1$, all the predicted ρ_{tr} are larger than $2 \times 10^7 \text{ gm cm}^{-3}$ by at least a factor of 2 except the case with $U = 10^6 \text{ cm/s}$. The predicted transition density with $K = 8$ is 2 – 3 times smaller than from $K = 1$. From Table 2, the predicted ρ_{tr} for the intermittency models with $K = 8$ agree with the observations within a factor of 2 except the case with a large velocity scale $U = 10^8 \text{ cm/s}$ at the integral length scale. To avoid discrepancy with the observations, our result indicates several possibilities.

1. The large scale motions caused by Rayleigh-Taylor instability freeze out due to the overall expansion of the star (Khokhlov 1995). The freezeout effect has to be efficient enough so that the developed part of the flow has a velocity scale of $\lesssim 10^6$ cm/s (see Table 2).
2. The flame is very robust. To break the flame, the local Gibson scale has to be at least $K^3 = 512$ times smaller than the flame thickness. In this case, the predicted transition density is 2 – 3 times smaller than from $K = 1$.
3. There is not enough time for the buoyancy-driven turbulence to fully develop down to the critical size before the predicted density for the DDT by our intermittency models is attained, thus motions at scales below the critical size are either absent or non-intermittent.
4. The DDT does not occur immediately after a region of the critical size enters the distributed regime. It may take some time for turbulence to help mix the region and make it nearly isothermal. However, the time scale for turbulence to mix a region of the critical size in the distributed regime is very small, $\lesssim 10^{-2}$ s at densities larger than 3×10^7 g/cm³. It is unlikely that the density drops much in such a short timescale.
5. Having a large enough region entering the distributed regime is not a sufficient condition for detonation. As mentioned in the Introduction, there are several uncertainties in the simple model by KOW and NW assuming flame quenching, entering the distributed regime and the DDT all occur simultaneously.

Our result alludes to the possibility that the criterion by KOW and NW is too weak for the DDT, supporting the claim of Lisewski et al (2000) and Woosley (2007) that just entering the distributed regime is not sufficient for the DDT. We have shown that their criteria for the DDT are much stronger than just entering the distributed regime. We also studied the intermittency effect on their conditions for the DDT. We find that the existence of a region of size 10^6 cm with rms turbulent velocity of 10^8 cm/s required by Lisewski et al (2007) for a DDT is rare, consistent with numerical results of Röpke (2007). We have also examined the intermittency effect on the transition density for the DDT criterion by Woosley (2007). We find that the effect is weaker for the stronger criterion and does not increase ρ_{tr} to above 10^7 g/cm³. Woosley (2007) obtained ρ_{tr} around 10^7 g/cm³ because of the assumption of strong turbulence velocity of 10^8 cm/s in a region of size 10^6 cm. Since the existence of such a region is rare, Woosley (2007) may considerably overestimate the transition density. This may imply that the condition for DDT by Lisewski et al. (2000) and Woosley (2007) is too strong and predicts a transition density smaller than that empirically determined from observations.

We have only studied the DDT in white dwarfs with the initial chemical composition of half carbon and half oxygen. In a white dwarf with more carbon, the nuclear timescale is smaller thus a stronger turbulent intensity is needed to break the flame. This results in a smaller transition density using the mean criterion (NW). On the other hand, the critical size for detonation in such a white dwarf is smaller, therefore the intermittency effect could be more efficient in increasing the transition density. We were not able to perform a calculation for the chemical composition with the carbon abundance larger than 0.5 due to the lack of sufficient data for the critical length scale in this case.

We point out that the intermittency models we used were originally proposed for homogeneous and isotropic turbulence. Turbulence in SNe Ia is stratified and may not achieve homogeneity and isotropy at very small scales even if the turbulence is developed at these scales. The effect of the departure from homogeneity and isotropy on the predicted transition density is out of the scope of this paper.

We thank Elaine Oran for useful discussions. This research was supported in part by NSF Grant AST-0707769 (LP, JCW) and by NASA Astrophysics Theory Program Grant NAG5-13280 (JS).

A. The She-Leveque Model

In their original paper, She and Leveque (1994) start with the moments of the distribution $P(\epsilon_l)$ and use the ratios of two successive moments, $\epsilon_l^{(p)} = \langle \epsilon_l^{p+1} \rangle / \langle \epsilon_l^p \rangle \propto l^{\tau_{p+1} - \tau_p}$, to characterize a hierarchy of dissipative structures. This ratio can be written as $\epsilon_l^{(p)} = \int \epsilon_l Q_p(\epsilon_l) d\epsilon_l$ where $Q_p(\epsilon_l) = \epsilon_l^p P(\epsilon_l) / \int \epsilon_l^p P(\epsilon_l) d\epsilon_l$. For a typical distribution $P(\epsilon_l)$ that decreases monotonically and faster than any power law at large ϵ_l , $Q_p(\epsilon_l)$ strongly peaks around $\epsilon_l^{(p)}$ for large p . Clearly $\epsilon_l^{(p)}$ increases with p , and $\epsilon_l^{(\infty)} = \lim_{n \rightarrow \infty} \langle \epsilon_l^{p+1} \rangle / \langle \epsilon_l^p \rangle$ corresponds to the most intense dissipative structures at scale l . These strongest dissipative structures are the origin of the anomalous scaling and the scaling of $\epsilon_l^{(\infty)}$ with l ,

$$\epsilon_l^{(\infty)} \sim l^{-\gamma} \quad (\text{A1})$$

is of fundamental importance. To determine the parameter γ , we can dimensionally write $\epsilon_l^{(\infty)}$ as an energy scale divided by a time scale t_l . She and Leveque argued that for the most intermittent structures this energy scale is the largest available kinetic energy (which is $\sim v_{rms}^2$, independent of l) and assumed that t_l exhibits a regular Kolmogorov scaling $t_l \sim l^{2/3}$, therefore $\gamma = 2/3$. From eq (A1), we have $\tau_{p+1} = \tau_p - \gamma$ for $p \rightarrow \infty$ or

$$\tau_p = -\gamma p + C, \quad p \rightarrow \infty \quad (\text{A2})$$

where the constant C has a physical interpretation as the codimension of the most intermittent structures. Eq (A1) means that the dissipation rate in a region of size l that encloses the most intensive structures scales with l as $\epsilon_l \propto l^{-\gamma}$. When calculating $\langle \epsilon_l^p \rangle$ at $p \rightarrow \infty$, we need to consider the possibility of a point finding itself within a distance l to the most intermittent structures, which is proportional to $l^{(D-d)}$ where D is the dimension of the system and d is the dimension of the most intensive structures (Frisch 1995). As $p \rightarrow \infty$, the contribution to $\langle \epsilon_l^p \rangle$ is dominated by the most intermittent structures, therefore $\langle \epsilon_l^p \rangle \propto l^{-\gamma p + D - d}$. Comparing with eq (A2), we find that C corresponds to the codimension of the most intermittent structures, $C = D - d$.

In order to determine the entire hierarchy of the dissipative structures, She and Leveque argued that the intensity $\epsilon_l^{(p+1)}$ of the dissipative structures at level $p + 1$ depends only on their immediate precursor, the structures of level p , from which the level $p + 1$ structures directly develop, and on the most intensive structures, where the structures of all orders tend to end up. Based on this argument, they made an assumption about the hierarchy of the dissipation rates,

$$\epsilon_l^{(p+1)} = A_p (\epsilon_l^{(p)})^\beta (\epsilon_l^{(\infty)})^{1-\beta} \quad (\text{A3})$$

where the coefficients A_p are independent of l but may be flow-dependent and non-universal. The parameter β will be completely fixed by γ and the codimension C . According to She and Leveque, this relation corresponds to a mysterious symmetry of the Navier-Stokes equation, termed "the hidden symmetry".

To derive τ_p from equation (A3), it is convenient to define a new variable,

$$\pi_l = \epsilon_l / \epsilon_l^{(\infty)} \quad (\text{A4})$$

which was introduced by Dubrulle (1994). Clearly $\epsilon_l^{(p)} / \epsilon_l^{(\infty)} = \langle \pi_l^{p+1} \rangle / \langle \pi_l^p \rangle$ from the definition of $\epsilon_l^{(p)}$. Then the "hidden symmetry" assumed by She and Leveque becomes,

$$\frac{\langle \pi_l^{p+2} \rangle}{\langle \pi_l^{p+1} \rangle} = A_p \left(\frac{\langle \pi_l^{p+1} \rangle}{\langle \pi_l^p \rangle} \right)^\beta \quad (\text{A5})$$

This recursion relation is solved by,

$$\frac{\langle \pi_l^{p+1} \rangle}{\langle \pi_l^p \rangle} = C_p \langle \pi_l \rangle^{\beta p} \quad (\text{A6})$$

where $C_0 = 1$ and $C_p = \prod_{n=0}^p A_n^{\beta n}$ for $p > 0$. Eq. (A6) gives,

$$\langle \pi_l^p \rangle = B_p \langle \pi_l \rangle^{(1-\beta p)/1-\beta} \quad (\text{A7})$$

where $B_0 = B_1 = 1$ and $B_p = \prod_{n=1}^{p-1} C_n$ for $p > 1$. Noting that $\langle \pi_l \rangle = \bar{\epsilon}/\epsilon_l^{(\infty)} \propto (\epsilon_l^{(\infty)})^{-1}$ and $\langle \pi_l^p \rangle = \langle \epsilon_l^p \rangle (\epsilon_l^{(\infty)})^{-p}$, eq (A7) gives $\langle \epsilon_l^p \rangle \propto (\epsilon_l^{(\infty)})^{p-(1-\beta^p)/(1-\beta)}$. Using eq (A1), we have

$$\tau_p = -\gamma p + \gamma \frac{(1 - \beta^p)}{1 - \beta} \quad (\text{A8})$$

The parameter β is determined by the asymptotic behavior of τ_p at the $p \rightarrow \infty$, eq (A2). Letting $p \rightarrow \infty$ in eq (A8) and comparing with eq (A2), we find that $\gamma/(1 - \beta) = C = D - d$. Since the most intermittent structures in 3D incompressible turbulence are filamentary, we have $d = 1$ and $C = 2$, thus $\beta = 1 - \gamma/2 = 2/3$ for $\gamma = 2/3$. Finally we arrive at the celebrated She-Leveque formulae,

$$\tau_p = -2p/3 + 2(1 - (2/3)^p) \quad (\text{A9})$$

and

$$\zeta_p = p/9 + 2(1 - (2/3)^{p/3}) \quad (\text{A10})$$

which agrees with the experimental result with an accuracy of 1%. Note that this result is consistent with the Kolmogorov's exact result for the third order structure function, i.e., $\zeta_3 = 1$. If the most intense structures are two dimensional, e.g., the dissipation sheets in compressible flows (Boldyrev et al. 2002) or the current sheets in MHD (Muller & Biskamp 2000), $\beta = 1/3$.

REFERENCES

- Anselmet, F., Gagne, Y., Hopfinger, E. J. & Antonia, R. A. 1984, J. Fluid Mech. 140, 63.
- Biskamp, D. 2003, Magnetohydrodynamic Turbulence. (Cambridge University Press)
- Boldyrev, S., Nordlund, A. & Padoan, P. 2002, Phys. Rev. Lett., 89, 031102
- Branch, D., Buta, R., Falk, S. W., McCall, M. L., Uomoto, A., Wheeler, J. C., Wills, B. J. & Sutherland, P. G. 1982, ApJ, 252, 61
- Branch, D., Lacy, C. H., McCall, M. L., Sutherland, P. G., Uomoto, A., Wheeler, J. C. & Wills, B. 1983, ApJ, 270, 123
- Dominguez, I., Höflich, P. & Straniero, O. 2001, ApJ, 557, 279.
- Dubrulle, B. 1994, Phys. Rev. Lett., 73, 959

- Frisch, U. 1995, *Turbulence*. (Cambridge University Press)
- Gamezo, V. N., Khokhlov, A. M., Oran, E. S., Chtchelkanova, A. Y., Rosenberg, R. O. 2003, *Science*, 299, 77
- Höflich, P. 1995, *ApJ*, 443, 89
- Höflich, P., Khokhlov, A. M., Wheeler, J. C. 1995, *ApJ*, 444, 83
- Höflich, P., Khokhlov 1996, *ApJ*, 457, 500
- Khokhlov, A. M. 1991, *A&A*, 245, 114
- Khokhlov, A. M. 1995, *ApJ*, 449, 695
- Khokhlov, A. M., Oran, E.S. & Wheeler, J. C. 1997 *ApJ*, 478, 678
- Kolmogorov, A. N. 1962, *J. Fluid Mech.* 13, 82
- Lisewski, A. M., Hillebrandt, W & Woosley, S. E. 2000, *ApJ*, 538, 831
- Lisewski, A. M., Hillebrandt, W., Woosley, S. E., Niemeyer, J. C. & Kerstein, A. R. 2000, *ApJ*, 537, 405 (Lisewski et al. 2000b)
- Monin, A. S. & Yaglom, A. M. 1975, *Statistical Fluid Mechanics*. (MIT press)
- Muller, W-C. & Biskamp, D 2000, *Phys. Rev. Lett.*, 84, 475
- Niemeyer, J. C. & Woosley, S. E. 1997 *ApJ*, 475, 740
- Niemeyer, J. C. & Kerstein 1997, *New Astronomy* 2, 239
- Oboukhov, A. M. 1962, *J. Fluid Mech.* 13, 77
- Röpke, F. K. 2007, *ApJ*, 668, 1103
- Röpke, F. K. & Hillebrandt, W. 2005, *A&A*, 431, 635
- She, Z-S. & Leveque, E. 1994, *Phys. Rev. Lett*, 72, 336
- She, Z-S. & Waymire, E. C. 1995, *Phys. Rev. Lett*, 74, 262
- Timmes, F. X. & Woosley, S. E. 1992, *ApJ*, 396, 649
- Wang, L-P., Chen, S., Brasseur, J. G. & Wyngaard, J. C. 1996, *J. Fluid Mech.* 309, 113
- Woosley, S. E. 2007, *ApJ*, 668, 1109

Yaglom, A. M. 1966, Dokl. Akad. Nauk SSSR, 166, 49

Yeung, P. K., Pope, S. B., Lamorgese, A. G. & Donzis, D. A. 2006, Phys. Fluids. 18, 065103

Zel’dovich, Y. B., Librovich, V. B., Makhviladze, G. M. & Sivashinsky, G. I. 1970, Acta Astron., 15., 313

Table 1: The laminar flame speed, the flame thickness and the critical length for a white dwarf with half carbon and half oxygen

ρ (10^9 g/cm ³)	S_l (10^5 cm/s)	l_f (cm)	ϵ_f (10^{15} cm ² /s ³)	l_c (cm)
2	75.8	9.35(-5)	4.66(9)	7(1)
0.5	18.1	9.46(-4)	6.27(6)	—
0.1	2.33	2.75 (-2)	4.60(2)	2(2)
0.05	0.599	5.19 (-1)	0.414	1.3(3) ^a
0.03	0.26 ^b	1.78 ^b	0.98 (-2)	5(3)
0.01	4.72(-2)	4.22	2.59(-5)	2(5)

^aRead from Fig. 6 in KOW.

^bRead from Fig. 7 in KOW.

Note. — The values of S_l and l_f are mainly taken from Table 3 of Timmes and Woosley 1992. The value of l_c is mainly taken from NW. We also include numbers (marked) from KOW because their results are very similar to *NW* despite the difference of details in the two models. Numbers in parentheses are powers of 10.

Table 2: The predicted transition densities ρ_{tr} (in unit of 10^7 g/cm³) for various models

Cases ^a	A	B	C	D	E
mean criterion	6.9(3.3)	5.5(2.3)	4.1(1.5)	3.0(1.0)	2.0 (—)
log- normal ^b	27(10)	11(4.9)	10(4.4)	9.4 (4.0)	4.3 (—)
log- normal ^c	23(7.6)	9.7(4.9)	8.4(3.9)	7.1 (3.0)	4.3 (—)
log- Poisson	24(8.6)	9.5(4.3)	8.7(3.8)	7.9(3.3)	3.8 (—)

^aA: $U = 10^8$ cm/s, $L = 10^8$ cm, $\bar{\epsilon} = 10^{16}$ cm²/s³;
B: $U = 10^7$ cm/s, $L = 10^6$ cm, $\bar{\epsilon} = 10^{15}$ cm²/s³;
C: $U = 10^7$ cm/s, $L = 10^7$ cm, $\bar{\epsilon} = 10^{14}$ cm²/s³;
D: $U = 10^7$ cm/s, $L = 10^8$ cm, $\bar{\epsilon} = 10^{13}$ cm²/s³;
E: $U = 10^6$ cm/s, $L = 10^6$ cm, $\bar{\epsilon} = 10^{12}$ cm²/s³

^bThe predicted transition density assuming a perfect fit of the distribution $P(\epsilon_{lc})$ by log-normal. This is the upper limit for ρ_{tr} since the log-normal approximation may break down and overestimate the distribution at the far tail.

^cThe lower limit for the transition density assuming the log-normal approximation applies only up to $5 - \sigma$.

Note. — The numbers in parentheses are the results predicted if the Gibson scale has to be 512 times smaller than the flame thickness for the transition to the distributed regime. In case E, (—) indicates that the transition density is smaller than 10^7 g/cm³, which cannot be well estimated since we only have data to 10^7 g/cm³ in Table 1.